

Topics in Computer Graphics

Chap 14: Tensor Product Patches

spring, 2014

University of Seoul
School of Computer Science
Minho Kim

Table of contents

Bilinear Interpolation

The Direct de Casteljaou Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Bilinear Interpolation

- ▶ Let $\mathbf{b}_{0,0}, \mathbf{b}_{0,1}, \mathbf{b}_{1,0}, \mathbf{b}_{1,1} \in \mathbb{E}^3$.
- ▶ The set of all points $\mathbf{x} \in \mathbb{E}^3$ of the form

$$\mathbf{x}(u, v) = \sum_{i=0}^1 \sum_{j=0}^1 \mathbf{b}_{i,j} B_i^1(u) B_j^1(v)$$

is called a *hyperbolic paraboloid* through the four points $\{\mathbf{b}_{i,j}\}$.

- ▶ In matrix form:

$$\mathbf{x}(u, v) = \begin{bmatrix} 1 - u & u \end{bmatrix} \begin{bmatrix} \mathbf{b}_{00} & \mathbf{b}_{01} \\ \mathbf{b}_{10} & \mathbf{b}_{11} \end{bmatrix} \begin{bmatrix} 1 - v \\ v \end{bmatrix}.$$

- ▶ $\mathbf{x}(u, v)$ is linear in both u and v
→ \mathbf{x} is called the *bilinear interpolant*.

Bilinear Interpolation (cont'd)

- ▶ Can be viewed as a map of the unit square $0 \leq u, v \leq 1$ in the u, v -plane. \rightarrow *domain*
- ▶ A line parallel to one of the axes in the domain corresponds to a curve in the range. (Equivalent to fixing either u or v .)
 \rightarrow *isoparametric curve*
- ▶ Every isoparametric curve of the hyperbolic paraboloid is a straight line.
 \rightarrow *ruled surface*
- ▶ Evaluation

$$\mathbf{b}_{0,0}^{0,1} = (1 - v)\mathbf{b}_{0,0} + v\mathbf{b}_{0,1}$$

$$\mathbf{b}_{1,0}^{0,1} = (1 - v)\mathbf{b}_{1,0} + v\mathbf{b}_{1,1}$$

$$\mathbf{x}(u, v) = \mathbf{b}_{0,0}^{1,1}(u, v) = (1 - u)\mathbf{b}_{0,0}^{0,1} + u\mathbf{b}_{1,0}^{0,1}$$

Equivalent to computing the coefficients of the isoparametric line $v = \text{const}$ first and then evaluating this

Bilinear Interpolation (cont'd)

- ▶ Why “hyperbolic paraboloid”?
 - ▶ The (nonparametric) surface $z = xy$ can be considered as the bilinear interpolant to the four points (Why?)

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

- ▶ The intersection of the surface and a plane parallel to the xy -plane is a hyperbola.
- ▶ The intersection of the surface and a plane containing the z -axis is a parabola.

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

The Direct de Casteljau Algorithm

- ▶ Higher degree surfaces can be obtained by repeated application of bilinear interpolation.
- ▶ Given points $\{\mathbf{b}_{i,j}\}_{i,j=0}^n$ and parameter values $(u, v) \in \mathbb{R}^2$, set

$$\mathbf{b}_{i,j}^{r,r} = [1 - u \quad u] \begin{bmatrix} \mathbf{b}_{i,j}^{r-1,r-1} & \mathbf{b}_{i,j+1}^{r-1,r-1} \\ \mathbf{b}_{i+1,j}^{r-1,r-1} & \mathbf{b}_{i+1,j+1}^{r-1,r-1} \end{bmatrix} \begin{bmatrix} 1 - v \\ v \end{bmatrix},$$

$i, j = 0, \dots, n - r$

and $\mathbf{b}_{i,j}^{0,0} = \mathbf{b}_{i,j}$. Then $\mathbf{b}_{0,0}^{n,n}(u, v)$ is the point with parameter values (u, v) on the *Bézier surface* $\mathbf{b}^{n,n}$.

- ▶ $\{\mathbf{b}_{i,j}\}$: Bézier points or control points
- ▶ The net of $\{\mathbf{b}_{i,j}\}$: Bézier net or control net of the surface $\mathbf{b}^{n,n}$
- ▶ What if the degrees in u and v are different?
($\{\mathbf{b}_{i,j}\}_{i=0,\dots,m,j=0,\dots,n}$)

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

The Tensor Product Approach

- ▶ “A surface is the locus of a curve that is moving through space and thereby changing its shape.”
- ▶ Formulation
 1. Given the initial Bézier curve of degree m :

$$\mathbf{b}^m(u) = \sum_{i=0}^m \mathbf{b}_i B_i^m(u)$$

2. Let each control point \mathbf{b}_i traverses a Bézier curve of degree n :

$$\mathbf{b}_i = \mathbf{b}_i(v) = \sum_{j=0}^n \mathbf{b}_{i,j} B_j^n(v).$$

3. Combining these two, we obtain the point $\mathbf{b}^{m,n}(u, v)$ on the surface $\mathbf{b}^{m,n}$ as

$$\mathbf{b}^{m,n}(u, v) = \sum_{i=0}^m \sum_{j=0}^n \mathbf{b}_{i,j} B_i^m(u) B_j^n(v).$$

The Tensor Product Approach (cont'd)

- ▶ For a fixed $\hat{v} = \text{const}$, its Bézier points are obtained by

$$\mathbf{b}_{i,0}^{0,n}(\hat{v}) = \sum_{j=0}^n \mathbf{b}_{ij} B_j^n(\hat{v}), \quad i = 0, \dots, m.$$

- ▶ Other straight lines in the domain (i.e. diagonals) are mapped to higher-degree (generally $m + n$) curves on the patch.

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Properties

- ▶ Affine invariance

$$\sum_{j=0}^n \sum_{i=0}^m B_i^m(u) B_j^n(v) \equiv 1(u, v)$$

→ $\mathbf{b}^{m,n}(u, v)$ is an affine combination of the control points.

- ▶ Convex hull property
- ▶ Boundary curves of the patch $\mathbf{b}^{m,n}(u, v)$ are polynomial curves.
- ▶ Variation diminishing property → NOT inherited

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Degree Elevation

- ▶ To rewrite a Bézier surface of degree (m, n) as one of degree $(m + 1, n)$:

$$\mathbf{b}^{m,n}(u, v) = \sum_{j=0}^n \left[\sum_{i=0}^{m+1} \mathbf{b}_{i,j}^{(1,0)} B_i^{m+1}(u) \right] B_j^n(v).$$

$$\mathbf{b}_{i,j}^{(1,0)} = \frac{i}{m+1} \mathbf{b}_{i-1,j} + \left(1 - \frac{i}{m+1}\right) \mathbf{b}_{i,j}, \quad \begin{cases} i = 0, \dots, m+1 \\ j = 0, \dots, n. \end{cases}$$

- ▶ as one of degree $(m + 1, n + 1)$:

$$\mathbf{b}_{i,j}^{(1,1)} = \begin{bmatrix} \frac{i}{m+1} & 1 - \frac{i}{m+1} \end{bmatrix} \begin{bmatrix} \mathbf{b}_{i-1,j-1} & \mathbf{b}_{i-1,j} \\ \mathbf{b}_{i,j-1} & \mathbf{b}_{i,j} \end{bmatrix} \begin{bmatrix} \frac{j}{n+1} \\ 1 - \frac{j}{n+1} \end{bmatrix}$$
$$\begin{cases} i = 0, \dots, m+1, \\ j = 0, \dots, n+1. \end{cases}$$

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Derivatives

- ▶ Partial derivatives along u and v .
- ▶ Tangent vector of an isoparametric curve.

$$\begin{aligned}\frac{\partial}{\partial u} \mathbf{b}^{m,n}(u, v) &= \sum_{j=0}^n \left[\frac{\partial}{\partial u} \sum_{i=0}^m \mathbf{b}_{i,j} B_i^m(u) \right] B_j^n(v) \\ &= m \sum_{j=0}^n \sum_{i=0}^m \Delta^{(1,0)} \mathbf{b}_{i,j} B_i^{m-1}(u) B_j^n(v).\end{aligned}$$

where $\Delta^{(1,0)} \mathbf{b}_{i,j} := \mathbf{b}_{i+1,j} - \mathbf{b}_{i,j}$.

$$\frac{\partial}{\partial v} \mathbf{b}^{m,n}(u, v) = n \sum_{i=0}^m \sum_{j=0}^n \Delta^{(0,1)} \mathbf{b}_{i,j} B_j^{n-1}(u) B_i^m(u).$$

Higher Derivatives

$$\frac{\partial^r}{\partial u^r} \mathbf{b}^{m,n}(u, v) = \frac{m!}{(m-r)!} \sum_{j=0}^n \sum_{i=0}^{m-r} \Delta^{(r,0)} \mathbf{b}_{i,j} B_i^{m-r}(u) B_j^n(v)$$

where $\Delta^{(r,0)} \mathbf{b}_{i,j} := \Delta^{(r-1,0)} \mathbf{b}_{i+1,j} - \Delta^{(r-1,0)} \mathbf{b}_{i,j}$ and

$$\frac{\partial^s}{\partial v^s} \mathbf{b}^{m,n}(u, v) = \frac{n!}{(n-s)!} \sum_{i=0}^m \sum_{j=0}^{n-s} \Delta^{(0,s)} \mathbf{b}_{i,j} B_j^{n-s}(v) B_i^m(u).$$

where $\Delta^{(0,s)} \mathbf{b}_{i,j} := \Delta^{(0,s-1)} \mathbf{b}_{i,j+1} - \Delta^{(0,s-1)} \mathbf{b}_{i,j}$.

Mixed Partial Derivatives & Cross Boundary Derivatives

- ▶ Mixed partial derivatives

$$\frac{\partial^{r+s}}{\partial u^r \partial v^s} \mathbf{b}^{m,n}(u, v) = \frac{m!n!}{(m-r)!(n-s)!} \sum_{i=0}^{m-r} \sum_{j=0}^{n-s} \Delta^{r,s} \mathbf{b}_{i,j} B_i^{m-r}(u) B_j^{n-s}(v).$$

- ▶ Cross boundary derivatives

$$\frac{\partial^r}{\partial u^r} \mathbf{b}^{m,n}(0, v) = \frac{m!}{(m-r)!} \sum_{j=0}^n \Delta^{(r,0)} \mathbf{b}_{0,j} B_j^n(0).$$

→ Important when we formulate condition for C^r continuity between adjacent patches.

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Blossoms

$$\mathbf{b}[u_1, \dots, u_m | v_1, \dots, v_n]$$

1. Compute the (curve) blossom value $\mathbf{b}_i[u_1, \dots, u_m]$ of all rows of control points, using the same values for each row.
2. Then use those values as input to the (curve) blossom $\mathbf{b}[v_1, \dots, v_n]$.
 - ▶ $\mathbf{b}[u^{<m>} | v^{<n>}]$ is the point on the surface.
 - ▶ The order of evaluation does not matter.
 - ▶ Multiaffinity in both u and v .

Outline

Bilinear Interpolation

The Direct de Casteljaou Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Curves on a Surface

1. Let $\mathbf{p} = (\mathbf{p}_u, \mathbf{p}_v)$ and $\mathbf{q} = (\mathbf{q}_u, \mathbf{q}_v)$ be u, v -coordinates of two points \mathbf{p} and \mathbf{q} in the domain of a tensor product patch of degree (n, n) .
2. A line through \mathbf{p} and \mathbf{q} :

$$\mathbf{u}(t) = (1 - t)\mathbf{p} + t\mathbf{q}.$$

3. A point on the curve is given by

$$\mathbf{b}[\left((1 - t)\mathbf{p}_u + t\mathbf{q}_u\right)^{\langle n \rangle} | \left((1 - t)\mathbf{q}_v + t\mathbf{q}_v\right)^{\langle n \rangle}].$$

4. Applying the Leibniz formula to the u -part,

$$\sum_{i+j=n} \binom{n}{i, j} (1 - t)^i t^j \mathbf{b}[\mathbf{p}_u^{\langle i \rangle} \mathbf{q}_u^{\langle j \rangle} | \left((1 - t)\mathbf{p}_v + t\mathbf{q}_v\right)^{\langle n \rangle}]$$

Curves on a Surface (cont'd)

5. Applying the Leibniz formula to the v -part,

$$\sum_{i+j=n} \sum_{r+s=n} \binom{n}{i, j} \binom{n}{r, s} (1-t)^i t^j (1-t)^r t^s \mathbf{b}[\mathbf{p}_u^{\langle i \rangle}, \mathbf{q}_u^{\langle j \rangle} | \mathbf{p}_v^{\langle r \rangle}, \mathbf{q}_v^{\langle s \rangle}]$$

6. Equivalently,

$$\begin{aligned} & \sum_{i=0}^n \sum_{j=0}^n \binom{n}{i} \binom{n}{j} (1-t)^{2n-i-j} t^{i+j} \mathbf{b}[\mathbf{p}_u^{\langle n-i \rangle}, \mathbf{q}_u^{\langle i \rangle} | \mathbf{p}_v^{\langle n-j \rangle}, \mathbf{q}_v^{\langle j \rangle}] \\ &= \sum_{k=0}^{2n} \sum_{i+j=k} \frac{\binom{n}{i} \binom{n}{j}}{\binom{2n}{k}} B_k^{2n}(t) \mathbf{b}[\mathbf{p}_u^{\langle n-i \rangle}, \mathbf{q}_u^{\langle i \rangle} | \mathbf{p}_v^{\langle n-j \rangle}, \mathbf{q}_v^{\langle j \rangle}] \\ &= \sum_{k=0}^{2n} \left(\sum_{i+j=k} \frac{\binom{n}{i} \binom{n}{j}}{\binom{2n}{k}} \mathbf{b}[\mathbf{p}_u^{\langle n-i \rangle}, \mathbf{q}_u^{\langle i \rangle} | \mathbf{p}_v^{\langle n-j \rangle}, \mathbf{q}_v^{\langle j \rangle}] \right) B_k^{2n}(t) \\ &=: \sum_{k=0}^{2n} \mathbf{c}_k B_k^{2n}(t) \end{aligned}$$

→ A curve of degree $2n$

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Normal Vectors

- ▶ Required for lighting computation.
- ▶ Can be obtained by cross product of any two tangent vectors at the given point on the surface.

$$\mathbf{b}(u, v) = \frac{\frac{\partial}{\partial u} \mathbf{b}^{m,n}(u, v) \wedge \frac{\partial}{\partial v} \mathbf{b}^{m,n}(u, v)}{\left\| \frac{\partial}{\partial u} \mathbf{b}^{m,n}(u, v) \wedge \frac{\partial}{\partial v} \mathbf{b}^{m,n}(u, v) \right\|}$$

- ▶ Degenerate cases?

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches

The Matrix Form of a Bézier patch

$$\mathbf{b}^{m,n}(u, v) = [B_0^m(u) \quad \cdots \quad B_m^m(u)] \begin{bmatrix} \mathbf{b}_{00} & \cdots & \mathbf{b}_{0n} \\ \vdots & \ddots & \vdots \\ \mathbf{b}_{n0} & \cdots & \mathbf{b}_{nn} \end{bmatrix} \begin{bmatrix} B_0^n(v) \\ \vdots \\ B_n^n(v) \end{bmatrix}$$

- ▶ $\{\mathbf{b}_{ij}\}$: Geometry matrix of the patch

Outline

Bilinear Interpolation

The Direct de Casteljau Algorithm

The Tensor Product Approach

Properties

Degree Elevation

Derivatives

Blossoms

Curves on a Surface

Normal Vectors

Twists

The Matrix Form of a Bézier patch

Nonparametric Patches